Author Index

Altmann, K.
— and Mukkur, T.K.S.
Alternative procedure for the purification of the heat-stable enterotoxin of enterotoxigenic *Escherichia coli* pathogenic for calves 253

Ang, S.B., see Lee, E.J.D. 203

Arai, K., see Koyama, I. 275

Avery, M.J.
— and Junk, G.A.
Procedure for the determination of amines in small urine samples 379

Bareggi, S.R., see Zecca, L. 417

Beijsterveldt, L.E.C. van, see Wuis, E.W. 212

Benchekroun, Y.
— , Ribon, B., Desage, M. and Brazier, J.L.
Simultaneous quantitation of phenobarbitone and *p*-hydroxyphenobarbitone and their perdeuteroethyl analogues in biological fluids by combined gas chromatography–mass spectrometry 287

Benet, L.Z., see Hyneck, M.L. 349

Beresford, A.P.
— , Macrae, P.V., Stopher, D.A. and Wood, B.A.
Analysis of amlodipine in human plasma by gas chromatography 178

Bhide, M.V., see Dunphy, M.J. 394

Björkman, S.
— and Elisson, L.O.
Determination of quinacrine (mepacrine) in plasma by high-performance liquid chromatography with fluorimetric detection 341

Blau, P.A.
— , Hines, J.W. and Voyksner, R.D.
Application of high-performance liquid chromatography and thermospray high-performance liquid chromatography–mass spectrometry to the analysis and identification of 2′,3′-dideoxyadenosine and its metabolite in biological media 1

Bochner, F., see Siebert, D.M. 425

Borghi, C., see Paroni, R. 189

Brazier, J.L., see Benchekroun, Y. 287

Brooks, P.M., see Dawson, M. 129

Bruijn, E.A. de, see Tjaden, U.R. 53

Carlier, J.-P.
— and Sellier, N.
Identification by gas chromatography–mass spectrometry of short-chain hydroxy acids produced by *Fusobacterium* species and *Clostridium innocuum* 121

Cazenave, J.-P., see Leray, C. 411

Chan, K., see Woo, J. 73

Chen, S.S., see Hsu, D.-S. 398

Choi, K.E.
— , Sinkule, J.A., Han, D.S., McGrath, S.C., Daly, K.M. and Larson, R.A.
High-performance liquid chromatographic assay for mitoxantrone in plasma using electrochemical detection 81

Dalgaard, L., see Hansen-Møller, J. 99

Daly, K.M., see Choi, K.E. 81

Dawson, M.
— , McGee, C.M., Brooks, P.M., Vine, J.H., Lacey, E. and Watson, T.R.
Quantitation of biphenylacetic acid in plasma and synovial fluid by gas chromatography–mass spectrometry–mass spectrometry 129

De Bruijn, E.A., see Tjaden, U.R. 53

Desage, M., see Benchekroun, Y. 287

Dirks, R.J.M., see Wuis, E.W. 212

Dobbs, N.A.
— and James, C.A.
Estimation of doxorubicin and doxorubicinol by high-performance liquid chromatography and advanced automated sample processor 184

Dunphy, M.J.
— , Bhide, M.V. and Smith, D.J.
Determination of hydroxyproline in tissue collagen hydrolysate by derivatization and isocratic reversed-phase high-performance liquid chromatography 394

Eckenhausen, M., see Powell, D.R. 163

Egger, H.-J.
— and Fischer, G.
Determination of the monocyclic β-lactam antibiotic carumonam in plasma and urine by ion-pair and ion-suppression reversed-phase high-performance liquid chromatography 357

Eggers, H., see Geschke, R. 111

Eichinger, A., see Nitsche, V. 207

Elisson, L.O., see Björkman, S. 341

Evelo, C.T.A.
— , Niessen, H.J.J.M., Roelofs, H.M.J. and Henderson, P.Th.
Cysteine adducts of human haemoglobin measured by isoelectric focusing in polyacrylamide gels with a non-linear pH gradient 35

Ferrario, P., see Zecca, L. 417
Fischer, G., see Egger, H.-J. 357
Fujita, K., see Imai, Y. 404
Gachályi, B., see Róna, K. 228
Geschke, R.
— , Körner, J. and Eggers, H.
Determination of the new monoamine oxidase inhibitor moclobemide and three of its metabolites in biological fluids by high-performance liquid chromatography 111
Greef, J. van der, see Tjaden, U.R. 53
Haginaka, J., see Nakagawa, T. 297
Han, D.S., see Choi, K.E. 81
Hansen, S.H., see Hansen-Møller, J. 99
Hansen-Møller, J.
— , Dalgaard, L. and Hansen, S.H.
Reversed-phase high-performance liquid chromatographic assay for the simultaneous determination of diflunisal and its glucuronides in serum and urine. Rearrangement of the 1-O-acylglucuronide 99
Hansson, C.
— , Kågedal, B. and Källberg, M.
Determination of 5-S-cysteinyldopa in human urine by direct injection in coupled-column high-performance liquid chromatography 146
Hartman, C.A., see Pivonka, J. 89
Hemmendinger, S., see Leray, C. 411
Henderson, P.Th., see Evelo, C.T.A. 35
Hines, J.W., see Blau, P.A. 1
Hintz, R.L., see Powell, D.R. 163
Hirai, Y., see Kitano, H. 13
Hoeven, R.A.M. van der, see Tjaden, U.R. 53
Hoppel, C.L., see Minkler, P.E. 385
Horitsu, H., see Idaka, E. 373
Hsu, D.-S.
— and Chen, S.S.
Assay of 5′-nucleotidase and simultaneous detection of interfering enzymes by anion-exchange column chromatography 398
Hyneck, M.L.
— , Smith, P.C. and Benet, L.Z.
High-performance liquid chromatographic determination of tolmetin, tolmetin glucuronide and its isomeric conjugates in plasma and urine 349
Hynninen, P.H., see Sainio, E.-L. 263
Idaka, E.
— , Kaiya, H. and Horitsu, H.
Determination of prostaglandin E_2 in plasma of schizophrenics and healthy controls by gas chromatography–negative-ion chemical ionization mass spectrometry and radioimmunoassay 373
Ikenata, T., see Omichi, K. 158
Ikezawa, H., see Koyama, I. 275
Imai, K., see Kotaniguchi, H. 141
Imai, Y.
— , Ito, S. and Fujita, K.
Determination of natural thiols by liquid chromatography after derivatization with 3,5-di-*tert.*-butyl-1,2-benzoquinone 404
Ingals, S.T., see Minkler, P.E. 385
Ise, N., see Kitano, H. 13
Ito, S., see Imai, Y. 404
James, C.A., see Dobbs, N.A. 184
Jol, C., see Tjaden, U.R. 53
Junk, G.A., see Avery, M.J. 379
Kågedal, B., see Hansson, C. 146
Kaiya, H., see Idaka, E. 373
Källberg, M., see Hansson, C. 146
Karlaganis, G.
— , Peretti, E. and Lautenburg, B.H.
Analysis of isoniazid, acetylhydrazine and [$^{15}N_2$]acetylhydrazine in serum by capillary gas chromatography–ammonia chemical ionization mass spectrometry 171
Kataoka, H.
— , Ohishi, K., Sumida, Y., Ohmori, M. and Makita, N.
Gas chromatographic assay for cysteine sulphinate decarboxylase activity in animal tissues 135
Kawakatsu, M., see Kotaniguchi, H. 141
Kawashima, T., see Nakagawa, T. 297
Kienle, M.G., see Paroni, R. 189
Kitano, H.
— , Hirai, Y., Okada, Y., Nakamura, K. and Ise, N.
Pulse injection analysis of the binding of serum proteins to porous polymer gels modified with formyl groups 13
Kleyn, E. van der, see Wuis, E.W. 212
Kobayashi, S.-I.
Simple method for the determination of nicardipine in plasma using high-performance liquid chromatography 439
Körner, J., see Geschke, R. 111
Kohse, K.P.
— and Wisser, H.
Quantitative monitoring by high-performance liquid chromatography of the dissociation of human serum cholinesterase by limited proteolysis 25
Komoda, T., see Koyama, I. 275
Kotaniguchi, H.
— , Kawakatsu, M., Toyo'oka, T. and Imai, K.
Automatic amino acid analysis utilizing 4-

fluoro-7-nitrobenzo-2-oxa-1,3-diazole 141
Koyama, I.
—, Arai, K., Sakagishi, Y., Ikezawa, H. and Komoda, T.
Blood appearance of rat alkaline phosphatase originating from the duodenum in vitro 275
Kucharczyk, N., see Pivonka, J. 89
Kuhnz, W.
Automated high-performance liquid chromatographic assay for cyproterone acetate and 15β-hydroxycyproterone acetate in plasma 432
Lacey, E., see Dawson, M. 129
Larking, P.W.
Quantitative gas chromatographic measurement of glycosaminoglycan hexosamines in urine and plasma 231
Larson, R.A., see Choi, K.E. 81
Lautenburg, B.H., see Karlaganis, G. 171
Lee, E.J.D.
—, Ang, S.B. and Lee, T.L.
Stereoselective high-performance liquid chromatographic assay for bupivacaine enantiomers 203
Lee, P.K.D., see Powell, D.R. 163
Lee, T.L., see Lee, E.J.D. 203
Leray, C.
—, Pelletier, X., Hemmendinger, S. and Cazenave, J.-P.
Thin-layer chromatography of human platelet phospholipids with fatty acid analysis 411
Lingeman, H., see Tjaden, U.R. 53
Macek, J.
— and Vácha, J.
Rapid and sensitive method for determination of piroxicam in human plasma by high-performance liquid chromatography 445
McGee, C.M., see Dawson, M. 129
McGrath, S.C., see Choi, K.E. 81
Macrae, P.V., see Beresford, A.P. 178
Makita, N., see Kataoka, H. 135
Maniwa, M., see Marunaka, T. 43, 329
Mårtensson, J.
Method for determination of free and total glutathione and γ-glutamylcysteine concentrations in human leukocytes and plasma 152
Marunaka, T.
—, Matsushima, E. and Maniwa, M.
Determination of cefodizime in biological materials by high-performance liquid chromatography 329
Marunaka, T.
—, Umeno, Y., Minami, Y., Matsushima, E., Maniwa, M., Yoshida, K. and Nagamachi, M.
Gas chromatographic–mass fragmentographic determination of propiverine and its metabolites in plasma and urine 43
Mascher, H.
High-performance liquid chromatographic determination of dextrorphan and 3-hydroxymorphinan in human plasma based on a selective pre-column sample clean-up 217
Matsushima, E., see Marunaka, T. 43, 329
Mefford, I.N.
—, Ota, M., Stipetic, M. and Singleton, W.
Application of a novel cation-exchange reagent, Igepon T-77 (N-methyl oleoyl taurate), to microbore separations of alumina extracts of catecholamines from cerebrospinal fluid, plasma, urine and brain tissue with amperometric detection 241
Minami, Y., see Marunaka, T. 43
Minkler, P.E.
—, Ingalls, S.T. and Hoppel, C.L.
Determination of total carnitine in human urine by high-performance liquid chromatography 385
Miyamoto, Y.
Highly sensitive determination of chlorpheniramine as fluorescence derivative by high-performance liquid chromatography 63
Mukkur, T.K.S., see Altmann, K. 253
Nagamachi, M., see Marunaka, T. 43
Nakagawa, T.
—, Shibukawa, A., Shimono, N., Kawashima, T., Tanaka, H. and Haginaka, J.
Retention properties of internal-surface reversed-phase silica packing and recovery of drugs from human plasma 297
Nakamura, K., see Kitano, H. 13
Niessen, H.J.J.M., see Evelo, C.T.A. 35
Nitsche, V.
—, Schütz, H. and Eichinger, A.
Rapid high-performance liquid chromatographic determination of nifedipine in plasma with on-line precolumn solid-phase extraction 207
Ohishi, K., see Kataoka, H. 135
Ohmori, M., see Kataoka, H. 135
Okada, Y., see Kitano, H. 13
Omichi, K.
— and Ikenaka, T.
Use of ion-exchange chromatography and

high-performance liquid chromatography to prepare a substrate for differential assay of human pancreatic and salivary α-amylases in serum 158
Ota, M., see Mefford, I.N. 241
Pang, K.S., see Xu, X. 313
Paroni, R.
— , Sirtori, C.R., Borghi, C. and Kienle, M.G.
High-performance liquid chromatographic determination of ribavirin in serum and urine and of its urinary metabolite 1,2,4-triazole-3-carboxamide 189
Pelletier, X., see Leray, C. 411
Peretti, E., see Karlaganis, G. 171
Pirola, R., see Zecca, L. 417
Pivonka, J.
— , Segelman, F.H., Hartman, C.A., Segl, W.E., Kucharczyk, N. and Sofia, R.D.
Determination of azelastine and desmethylazelastine in human plasma by high-performance liquid chromatography 89
Powell, D.R.
— , Lee, P.D.K., Shively, J.E., Eckenhausen, M. and Hintz, R.L.
Method for publication of an insulin-like growth factor-binding protein produced by human HEP G2 hepatoma cells 163
Puchta, V.
— and Spiteller, G.
Identification of 3-hydroxydecandioic acid dimethylate in human urine 450
Ribon, B., see Benchekroun, Y. 287
Roelofs, H.M.J., see Evelo, C.T.A. 35
Róna, K.
— and Gachályi, B.
Simple liquid chromatographic method for the determination of ornidazole and metronidazole in human serum 228
Sainio, E.-L.
— and Hynninen, P.H.
High-voltage electrophoretic separation of tryptophan metabolites of the kynurenine pathway 263
Sakagishi, Y., see Koyama, I. 275
Šalamoun, J.
— , Šprta, V., Sládek, T. and Smrž, M.
Determination of acyclovir in plasma by column liquid chromatography with fluorescence detection 197
Schütz, H., see Nitsche, V. 207
Segelman, F.H., see Pivonka, J. 89
Segl, W.E., see Pivonka, J. 89
Sellier, N., see Carlier, J.-P. 121
Shibukawa, A., see Nakagawa, T. 297
Shimono, N., see Nakagawa, T. 297
Shively, J.E., see Powell, D.R. 163
Siebert, D.M.
— and Bochner, F.
Determination of plasma aspirin and salicylic acid concentrations after low aspirin doses by high-performance liquid chromatography with post-column hydrolysis and fluorescence detection 425
Singleton, W., see Mefford, I.N. 241
Sinkule, J.A., see Choi, K.E. 81
Sirtori, C.R., see Paroni, R. 189
Sládek, T., see Šalamoun, J. 197
Smith, D.J., see Dunphy, M.J. 394
Smith, P.C., see Hyneck, M.L. 349
Smrž, M., see Šalamoun, J. 197
Sofia, R.D., see Pivonka, J. 89
Spiteller, G., see Puchta, V. 450
Šprta, V., see Šalamoun, J. 197
Stipetic, M., see Mefford, I.N. 241
Stopher, D.A., see Beresford, A.P. 178
Sumida, Y., see Kataoka, H. 135
Tanaka, H., see Nakagawa, T. 297
Teoh, R., see Woo, J. 73
Tjaden, U.R.
— , DeBruijn, E.A., Van der Hoeven, R.A.M., Jol, C., Van der Greef, J. and Lingeman, H.
Automated analysis of mitomycin C in body fluids by high-performance liquid chromatography with on-line sample pre-treatment 53
Toyo'oka, T., see Kotaniguchi, H. 141
Umeno, Y., see Marunaka, T. 43
Vácha, J., see Macek, J. 445
Van Beijsterveldt, L.E.C., see Wuis, E.W. 212
Van der Greef, J., see Tjaden, U.R. 53
Van der Hoeven, R.A.M., see Tjaden, U.R. 53
Van der Kleyn, E., see Wuis, E.W. 212
Vine, J.H., see Dawson, M. 129
Voyksner, R.D., see Blau, P.A. 1
Vree, T.B., see Wuis, E.W. 212
Ward, R.
— and Workman, P.
Gradient high-performance liquid chromatographic method for simultaneous assay of the radiosensitizers etanidazole (SR 2508) and pimonidazole (Ro 03-8799) in biological materials 223
Watson, T.R., see Dawson, M. 129
Wisser, H., see Kohse, K.P. 25
Wong, C.L., see Woo, J. 73
Woo, J.
— , Wong, C.L., Teoh, R. and Chan, K.
Liquid chromatographic assay for the

simultaneous determination of pyrazinamide and rifampicin in serum samples from patients with tuberculous meningitis 73
Wood, B.A., see Beresford, A.P. 178
Workman, P., see Ward, R. 223
Wuis, E.W.
— , Van Beijsterveldt, L.E.C., Dirks, R.J.M., Vree, T.B. and Van der Kleyn, E.
Rapid simultaneous determination of baclofen and its γ-hydroxy metabolite in urine by high-performance liquid chromatography with ultraviolet detection 212
Xu, X.
— and Pang, K.S.
High-performance liquid chromatographic method for the quantitation of salicylamide and its metabolites in biological fluids 313
Yoshida, K., see Marunaka, T. 43
Zecca, L.
— , Ferrario, P., Pirola, R. and Bareggi, S.R.
Analysis of chlorodesmethyldiazepam and its metabolites in plasma and urine 417

Subject Index

Acetylhydrazine
Analysis of isoniazid acetylhydrazine and [$^{15}N_2$]acetylhydrazine in serum by capillary GC–ammonia chemical ionization MS 171

Acetylsalicylic acid
Determination of plasma aspirin and salicylic acid concentrations after low aspirin doses by HPLC with post-column hydrolysis and fluorescence detection 425

Acyclovir
Determination of acyclovir in plasma by column LC with fluorescence detection 197

Alkaline phosphatase
Assay of 5′-nucleotidase and simultaneous detection of interfering enzymes by anion-exchange column chromatography 398

Alkaline phosphatase
Blood appearance of rat alkaline phosphatase originating from the duodenum in vitro 275

Amines
Application of a novel cation-exchange reagent, Igepon T-77 (N-methyl oleoyl taurate), to microbore separations of alumina extracts of catecholamines from cerebrospinal fluid, plasma, urine and brain tissue with amperometric detection 241

Amines
Procedure for the determination of amines in small urine samples 379

Amino acids
Automatic amino acid analysis utilizing 4-fluoro-7-nitrobenzo-2-oxa-1,3-diazole 141

Amlodipine
Analysis of amlodipine in human plasma by GC 178

α-Amylases
Use of ion-exchange chromatography and HPLC to prepare a substrate for differential assay of human pancreatic salivary α-amylases in serum 158

Anthranilic acid
High-voltage electrophoretic separation of tryptophan metabolites of the kynurenine pathway 263

Aspirin
Determination of plasma aspirin and salicylic acid concentrations after low aspirin doses by HPLC with post-column hydrolysis and fluorescence detection 425

Azelastine
Determination of azelastine and desmethylazelastine in human plasma by HPLC 89

Baclofen
Rapid simultaneous determination of baclofen and its γ-hydroxy metabolite in urine by HPLC with ultraviolet detection 212

Biphenylacetic acid
Quantitation of biphenylacetic acid in plasma and synovial fluid by GC-MS-MS 129

Bisantrene
HPLC assay for mitoxantrone in plasma using electrochemical detection 81

Bupivacaine
Stereoselective HPLC assay for bupivacaine enantiomers 203

Cardiolipin
TLC of human platelet phospholipids with fatty acid analysis 411

Carnitine
Determination of total carnitine in human urine by HPLC 385

Carumonam
Determination of the monocyclic β-lactam antibiotic carumonam in plasma and urine by ion-pair and ion-suppression reversed-phase HPLC 357

Catecholamines
Application of a novel cation-exchange reagent, Igepon T-77 (N-methyl oleoyl taurate), to microbore separations of alumina extracts of catecholamines from cerebrospinal fluid, plasma, urine and brain tissue with amperometric detection 241

Cefaclor
Retention properties of internal-surface reversed-phase silica packing and recovery of drugs from human plasma 297

Cefodizime
Determination of cefodizime in biological materials by HPLC 329

Cefpiramide
Retention properties of interal-surface reversed-phase silica packing and recovery of drugs from human plasma 297

Chlorodesmethyldiazepam
Analysis of chlorodesmethyldiazepam and its metabolites in plasma and urine 417

Chlorpheniramine
Highly sensitive determination of chlorpheniramine as fluorescence derivative by HPLC 63
Cholinesterase
Quantitative monitoring by HPLC of the dissociation of human serum cholinesterase by limited proteolysis 25
Clostridium innocuum
Identification by GC–MS of short-chain hydroxy acids produced by *Fusobacterium* species and *Clostridium innocuum* 121
Cyproterone acetate
Automated HPLC assay for cyproterone acetate and 15β-hydroxycyproterone acetate in plasma 432
L-Cysteine
Determination of natural thiols by LC after derivatization with 3,5-di-*tert.*-butyl-1,2-benzoquinone 404
Cysteine sulphinate decarboxylase
GC assay for cysteine sulphinate decarboxylase activity in animal tissues 135
5-S-Cysteinyldopa
Determination of 5-S-cysteinyldopa in human urine by direct injection in coupled-column HPLC 146
Desmethylazelastine
Determination of azelastine and desmethylazelastine in human plasma by HPLC 89
Dextrorphan
HPLC determination of dextrorphan and 3-hydroxymorphinan in human plasma based on a selective pre-column sample clean-up 217
3,5-Di-*tert.*-butyl-1,2-benzoquinone
Determination of natural thiols by LC after derivatization with 3,5-di-*tert.*-butyl-1,2-benzoquinone 404
2′,3′-Dideoxyadenosine
Application of HPLC and thermospray HPLC–MS to the analysis and identification of 2′,3′-dideoxyadenosine and its metabolite in biological media 1
2′,3′-Dideoxyinosine
Application of HPLC and thermospray HPLC–MS to the analysis and identification of 2′,3′-dideoxyadenosine and its metabolite in biological media 1
Diflunisal
Reversed-phase HPLC assay for the simultaneous determination of diflunisal and its glucuronides in serum and urine. Rearrangement of the 1-O-acetylglucuronide 99
Diflunisal ester glucuronide
Reversed-phase HPLC assay for the simultaneous determination of diflunisal and its glucuronides in serum and urine. Rearrangement of the 1-O-acetylglucuronide 99
Diflunisal ether glucuronide
Reversed-phase HPLC assay for the simultaneous determination of diflunisal and its glucuronides in serum and urine. Rearrangement of the 1-O-acetylglucuronide 99
Doxorubicin
Estimation of doxorubicin and doxorubicinol by HPLC and advanced automated sample processor 184
Doxorubicinol
Estimation of doxorubicin and doxorubicinol by HPLC and advanced automated sample processor 184
Enterotoxin
Alternative procedure for the purification of the heat-stable enterotoxin of enterotoxigenic *Escherichia coli* pathogenic for calves 253
Enzymes
Assay of 5′-nucleotidase and simultaneous detection of interfering enzymes by anion-exchange column chromatography 398
Enzymes
Blood appearance of rat alkaline phosphatase originating from the duodenum in vitro 275
Enzymes
GC assay for cysteine sulphinate decarboxylase activity in animal tissues 135
Enzymes
Quantitative monitoring by HPLC of the dissociation of human serum cholinesterase by limited proteolysis 25
Enzymes
Use of ion-exchange chromatography and HPLC to prepare a substrate for differential assay of human pancreatic and salivary α-amylases in sereum 158
Escherichia coli enterotoxin
Alternative procedure for the purification of the heat-stable enterotoxin of enterotoxigenic *Escherichia coli* pathogenic for calves 253
Etanidazole
Gradient HPLC method for simultaneous assay of the radiosensitizers etanidazole (SR 2508) and pimonidazole (Ro 03-8799) in biological materials 223
Fatty acid analysis
TLC of human platelet phospholipids with fatty acid analysis 411

4-Fluoro-7-nitrobenzo-2-oxa-1,3-diazole
Automatic amino acid analysis utilizing 4-fluoro-7-nitrobenzo-2-oxa-1,3-diazole 141
Fusobacterium species
Identification by GC–MS of short-chain hydroxy acids produced by *Fusobacterium* species and *Clostridium innocuum* 121
Gentisamide
HPLC method for the quantitation of salicylamide and its metabolites in biological fluids 313
Gentisamide 5-glucuronide
HPLC method for the quantitation of salicylamide and its metabolites in biological fluids 313
Gentisamide sulphates
HPLC method for the quantitation of salicylamide and its metabolites in biological fluids 313
γ-Glutamylcysteine
Method for determination of free and total glutathione and γ-glutamylcysteine concentrations in human leukocytes and plasma 152
Glutathione
Determination of natural thiols by LC after derivatization with 3,5-di-*tert.*-butyl-1,2-benzoquinone 404
Glutathione
Method for determination of free and total glutathione and γ-glutamylcysteine concentrations in human leukocytes and plasma 152
Glycosaminoglycan hexosamines
Quantitative GC measurement of glycosaminoglycan hexosamines in urine and plasma 231
Glycosaminoglycans
Quantitatie GC measurement of glycosaminoglycan hexosamines in urine and plasma 231
Haemoglobin, cysteine adducts
Cysteine adducts of human haemoglobin measured by isoelectric focusing in polyacrylamide gels with a non-linear pH gradient 35
HEP G2 hepatoma cells
Method for purification of an insulin-like growth factor-binding protein produced by human HEP G2 hepatoma cells 163
Hexosamines
Quantitative GC measurement of glycosaminoglycan hexosamines in urine and plasma 231
Hydroxy acids, short-chain
Identification by GC–MS of short-chain hydroxy acids produced by *Fusobacterium* species and *Clostridium innocuum* 121
Hydroxyanthranilic acid
High-voltage electrophoretic separation of tryptophan metabolites of the kynurenine pathway 263
γ-Hydroxybaclofen
Rapid simultaneous determination of baclofen and its γ-hydroxy metabolite in urine by HPLC with ultraviolet detection 212
15β-Hydroxycyproterone acetate
Automated HPLC assay for cyproterone acetate and 15β-hydroxycyproterone acetate in plasma 432
3-Hydroxydecandioic acid dimethylate
Identification of 3-hydroxydecandioic acid dimethylate in human urine 450
3-Hydroxymorphinan
HPLC determination of dextrorphan and 3-hydroxymorphinan in human plasma based on a selective pre-column sample clean-up 217
p-Hydroxyphenobarbitone
Simultaneous quantitation of phenobarbitone and *p*-hydroxyphenobarbitone and their perdeuteroethyl analogues in biological fluids by combined GC–MS 287
Hydroxyproline
Determination of hydroxyproline in tissue collagen hydrolysate by derivatization and isocratic reversed-phase HPLC 394
Igepon T-77
Application of a novel cation-exchange reagent, Igepon T-77 (N-methyl oleoyl taurate), to microbore separations of alumina extracts of catecholamines from cerebrospinal fluid, plasma, urine and brain tissue with amperometric detection 241
Insulin-like growth factor-binding protein
Method for purification of an insulin-like growth factor-binding protein produced by human HEP G2 hepatoma cells 163
Internal-surface reversed-phase silica
Retention properties of internal-surface reversed-phase silica packing and recovery of drugs from human plasma 297
Isoniazid
Analysis of isoniazid acetylhydrazine and [$^{15}N_2$]acetylhydrazine in serum by capillary GC–ammonia chemical ionization MS 171

Kynurenic acid
High-voltage electrophoretic separation of tryptophan metabolites of the kynurenine pathway 263
Kynurenine
High-voltage electrophoretic separation of tryptophan metabolites of the kynurenine pathway 263
Kynurenine pathway
High-voltage electrophoretic separation of tryptophan metabolites of the kynurenine pathway 263
β-Lactam antibiotics
Determination of the monocyclic β-lactam antibiotic carumonam in plasma and urine by ion-pair and ion-suppression reversed-phase HPLC 357
Lidocaine
Retention properties of internal-surface reversed-phase silica packing and recovery of drugs from human plasma 297
Lipids
TLC of human platelet phospholipids with fatty acid analysis 411
Lorazepam
Analysis of chlorodesmethyldiazepam and its metabolites in plasma and urine 417
Mepacrine
Determination of quinacrine (mepacrine) in plasma by HPLC with fluorimetric detection 341
N-Methyl oleoyl taurate
Application of a novel cation-exchange reagent, Igepon T-77 (N-methyl oleoyl taurate), to microbore separations of alumina extracts of catecholamines from cerebrospinal fluid, plasma, urine and brain tissue with amperometric detection 241
1-Methyl-4-piperidyl diphenyl propoxyacetate
GC–mass fragmentographic determination of propiverine and its metabolites in plasma and urine 43
Metronidazole
Simple LC method for the determination of ornidazole and metronidazole in human serum 228
Mitomycin C
Automated analysis of mitomycin C in body fluids by HPLC with on-line sample pretreatment 53
Mitoxantrone
HPLC assay for mitoxantrone in plasma using electrochemical detection 81
Moclobemide
Determination of the new monoamine oxidase inhibitor moclobemide and three of its metabolites in biological fluids by HPLC 111
Nicardipine
Simple method for the determination of nicardipine in plasma using HPLC 439
Nicotinic acid
High-voltage electrophoretic separation of tryptophan metabolites of the kynurenine pathway 263
Nifedipine
Rapid HPLC determination of nifedipine in plasma with on-line precolumn solid-phase extraction 207
Nucleosides
Application of HPLC and thermospray HPLC–MS to the analysis and identification of 2′,3′-dideoxyadenosine and its metabolite in biological media 1
5′-Nucleotidase
Assay of 5′-nucleotidase and simultaneous detection of interfering enzymes by anion-exchange column chromatography 398
Ornidazole
Simple LC method for the determination of ornidazole and metronidazole in human serum 228
Phenobarbitone
Simultaneous quantitation of phenobarbitone and *p*-hydroxyphenobarbitone and their perdeuteroethyl analogues in biological fluids by combined GC–MS 287
Phosphatidic acid
TLC of human platelet phospholipids with fatty acid analysis 411
Phosphatidylethanolamine
TLC of human platelet phospholipids with fatty acid analysis 411
Phosphatidylglycerol
TLC of human platelet phospholipids with fatty acid analysis 411
Phosphatidylinositol
TLC of human platelet phospholipids with fatty acid analysis 411
Phosphatidylserine
TLC of human platelet phospholipids with fatty acid analysis 411
Phospholipids
TLC of human platelet phospholipids with fatty acid analysis 411

Pimonidazole
Gradient HPLC method for simultaneous assay of the radiosensitizers etanidazole (SR 2508) and pimonidazole (Ro 03-8799) in biological materials 223
Piroxicam
Rapid and sensitive method for determination of piroxicam in human plasma by HPLC 445
Probenecid
Retention properties of internal-surface reversed-phase silica packing and recovery of drugs from human plasma 297
Propiverine
GC–mass fragmentographic determination of propiverine and its metabolites in plasma and urine 43
Prostaglandin E_2
Determination of prostaglandin E_2 in plasma of schizophrenics and healthy controls by GC–negative-ion chemical ionization MS and radioimmunoassay 373
Proteins
Cysteine adducts of human haemoglobin measured by isoelectric focusing in polyacrylamide gels with a non-linear pH gradient 35
Proteins
Method for purification of an insulin-like growth factor-binding protein produced by human HEP G2 hepatoma cells 163
Proteins
Pulse injection analysis of the binding of serum proteins to porous polymer gels modified with formyl groups 13
Pyrazinamide
LC assay for the simultaneous determination of pyrazinamide and rifampicin in serum samples from patients with tuberculous meningitis 73
Quinacrine
Determination of quinacrine (mepacrine) in plasma by HPLC with fluorimetric detection 341
Quinolinic acid
High-voltage electrophoretic separation of tryptophan metabolites of the kynurenine pathway 263
Ribavirin
HPLC determination of ribavirin in serum and urine and of its urinary metabolite 1,2,4-triazole-3-carboxamide 189
Rifampicin
LC assay for the simultaneous determination of pyrazinamide and rifampicin in serum samples from patients with tuberculous meningitis 73
Salicylamide
HPLC method for the quantitation of salicylamide and its metabolites in biological fluids 313
Salicylamide glucuronide
HPLC method for the quantitation of salicylamide and its metabolites in biological fluids 313
Salicylamide sulphate
HPLC method for the quantitation of salicylamide and its metabolites in biological fluids 313
Salicylic acid
Determination of plasma aspirin and salicylic acid concentrations after low aspirin doses by HPLC with post-column hydrolysis and fluorescence detection 425
Serum proteins
Pulse injection analysis of the binding of serum proteins to porous polymer gels modified with formyl groups 13
Stationary phases
Retention properties of internal-surface reversed-phase silica packing and recovery of drugs from human plasma 297
Thiols
Determination of natural thiols by LC after derivatization with 3,5-di-*tert.*-butyl-1,2-benzoquinone 404
Tolmetin
HPLC determination of tolmetin, tolmetin glucuronide and its isomeric conjugates in plasma and urine 349
Tolmetin glucuronide
HPLC determination of tolmetin, tolmetin glucuronide and its isomeric conjugates in plasma and urine 349
1,2,4-Triazole-3-carboxamide
HPLC determination of ribavirin in serum and urine and of its urinary metabolite 1,2,4-triazole-3-carboxamide 189
Tryptophan
High-voltage electrophoretic separation of tryptophan metabolites of the kynurenine pathway 263
Xanthurenic acid
High-voltage electrophoretic separation of tryptophan metabolites of the kynurenine pathway 263